arXiv:1304.5522v1 [gr-qc] 7 Apr 2013

Velocity requirements for causality violation

G. Modanese *[1]
[1]*Free University of Bolzano, Faculty of Science and Technology, Bolzano, Italy*

(Dated: April 6, 2013)

It is known that the hypothetical existence of superluminal signals would imply the logical possibility of active causal violation: an observer in relative motion with respect to a primary source could in principle emit secondary superluminal signals (triggered by the primary ones) which go back in time and deactivate the primary source before the initial emission. This is a direct consequence of the structure of the Lorentz transformations, sometimes called "Regge-Tolman paradox". It is straightforward to find a formula for the velocity of the moving observer required to produce the causality violation. When applied to some recent claims of slight superluminal propagation, this formula yields a required velocity very close to the speed of light; this raises some doubts about the real physical observability of such violations. We re-compute this velocity requirement introducing a realistic delay between the reception of the primary signal and the emission of the secondary. It turns out that for *any* delay it is possible to find moving observers able to produce active causal violation. This conclusion looks physically absurd and is mathematically due to the singularity of the Lorentz transformations for $\beta \to 1^-$. It appears to undermine the connection between superluminal signals and active causality violation. We also briefly consider further paradoxes raised by superluminal propagation: the contemporary emission of several superluminal signals from a source and the interaction of two particles through the exchange of superluminal virtual particles.

PACS numbers: 2.25.Bs, 03.30.+p

I. INTRODUCTION

Phenomena of electromagnetic wave propagation with superluminal group velocity have been observed in several laboratories in the last years and can be collected in two categories: (a) evanescent waves, and (b) Bessel beams of so-called "X-shaped waves".

Concerning the first category, superluminal effects for evanescent waves have been demonstrated in tunnelling experiments in both the optical domain and microwaves range [1]; these effects can be revealed, however, only over short distances, typically a few centimetres for microwaves (the most favourable case).

Concerning the second category, Mugnai et al. have demonstrated the superluminal propagation of localized microwaves over a distance of 1 m or more [2]. Let us summarize the experiment very briefly. A carrier signal of frequency 8.6 GHz was modulated with rectangular pulses; the waves were first sent by a horn antenna on a circular slit with mean diameter d placed in the focal plane of a circular mirror. In this way, a so-called Bessel beam was generated by interference. The field of the beam can be considered as formed by the superposition of pairs of X-shaped plane waves. These move along the axis of the mirror with velocity approximately 5-7% larger than the

---

* Email address: giovanni.modanese@unibz.it

light velocity, for a slit diameter d of 20 cm, or up to 25% for d=10 cm. A similar experiment was performed in the optical range [3], but a clear observation of superluminal propagation was impossible in that case.

Several papers ([2, 7, 8] and ref.s) discuss the issue of signal transmission in experiments of this kind. The question is, if superluminal propagation effects can be used to convey information at superluminal speed, and the answer is generally that they can't, though it also depends on what is exactly meant by a signal. A typical argument is that waves with superluminal group velocity are always accompanied by a "precursor wave" propagating at light speed. Some authors speculated, however, that in certain cases the superluminal wave could overtake the precursor. Other authors argued that the violation of causality by true superluminal signals is only apparent and could be avoided through the so-called Feynman-Stuckelberg tachyon reinterpretation principle.

Data from the OPERA collaboration [4] in 2011 pointed to the possible existence of superluminal neutrinos with velocity in excess of c by about 20 parts in one million. Later data by ICARUS and OPERA itself failed to confirm the result. At the astrophysical level, observation of neutrinos from the supernova SN1987A set an upper bound of $2 \cdot 10^{-9}$ on possible relative deviations from light velocity [5].

In this work we analyze some of the main reasons why superluminal propagation phenomena, if they exist, are thought to violate causality. We follow in part a pedagogical approach and stress the fundamental aspects, but we also obtain some new results. We draw a distinction between "passive" causality violation (the possibility to *observe* a cause-effect inversion in a suitable moving system) and "active" causality violation (the possibility that an effect triggers the disabling of its own cause). For both cases we compute the velocity that the moving system must have in order to make the causality violation possible (Sect.s II, III). In the case of active violation, we consider the possibility that there is a dead time in the re-emission of the superluminal signal (Sect. IV). We include in the calculation also extreme cases, with large superluminality and/or large delays. Finally, in the case of passive violation we consider an hypothetical source which emits several superluminal particles at the same time, in a sort of "explosive" process; we find that this process, as seen from a moving system, appears to violate the entropic principle. We also analyse the exchange of tachyons between two real particles from the point of view of a moving system (Sect. V). (It is known that quantum field theories with tachyons are plagued by instabilities; here we consider tachyons not as fundamental particles, but only as a possible formal representation of superluminal signals.)

The conclusions correspond only in part to the possible intuitive expectations. In particular:

(1) For small superluminality ($V = c(1+\varepsilon)$, $\varepsilon \ll 1$) the velocity v of the moving system required for both types of causality violation is very large and quite unrealistic. We have $v \simeq (1-\varepsilon)c$ for passive violation and $v \simeq (1-\varepsilon^2/2)c$ for active violation.

(2) In the presence of re-transmission delays the active violation is always possible, and this is paradoxical, because the delay can also be very large compared to the propagation time. This means, in our opinion, that in the presence of delay the active violation is an artifact of the Lorentz transformations, due to their singularity for $\beta \to 1^-$. Such a singularity could be eliminated through a physical cut-off, based for instance on the existence of a maximum acceleration for reference frames [9, 10]. Therefore the presence of re-transmission delay, which is physically unavoidable, eliminates in many cases the possibility of active violations and leaves only the possibility of passive violations. Passive violations, however, are in some sense much less pathological, and are actually typical of EPR-like phenomena which require a non-superluminal "precursor" for state preparation.

(3) Considering an exchange of tachyons between two real particles from the point of view of a moving system, we find this process to be inconsistent with local realism: if two bodies interact through the exchange of tachyons, it is impossible to say which body emits them and which one absorbs them. We can only consider the exchange process as a whole.

II. SUPERLUMINAL PROPAGATION VS. CAUSALITY VIOLATION. THE CASE OF "PASSIVE VIOLATION"

One often finds in the literature the statement that "the existence of a correlation between events separated by a space-like interval implies a violation of causality". (These are two events such that their spatial distance Δx is larger than $c\Delta t$.) The reason is that there exist suitable reference systems in relative motion where the temporal order of the events appears to be reversed, i.e. the delay $\Delta t'$ becomes negative. A legitimate question then arises: how fast must such relative motion be, for the moving observer to see the temporal order of the two events reversed? A simple exercise on Lorentz transformations gives the answer: if the two events appear to be connected by a superluminal signal with velocity V (i.e., $\Delta x/\Delta t = V > c$ in the rest reference system), then the causality violation is observed in reference systems moving with relative velocity v larger than c^2/V.

Namely, suppose that the superluminal signal is emitted at the origin ($x = 0, t = 0$) of a spacetime reference system. (Coordinates y and z are omitted, only boosts along x are considered.) The signal is received at a point $R(x,t)$ outside the light-cone (see Fig. 1). We have $t < x/c$, and more exactly $t = x/V$, where V is the signal velocity ($V > c$). All this is in a reference system



FIG. 1: The "light cone" (here seen in a 2D section) separates the points of spacetime which are in causal relationship to the point $(0,0)$ from those, like R, which are not.

where emitter and receiver are at rest.

Now make a Lorenz boost with parameters β, γ along x; the new coordinates in the "moving" system are $t' = \gamma(t - \beta x/c)$, $x' = \gamma(x - \beta ct)$. We look for values of β such that $t' < 0$, ie the event R is seen in the new reference system as antecedent to the emission. (Note that the emission remains at $(0,0)$ in the new coordinates.) Putting $t' = 0$ we find $t = \beta x/c$ and remembering that $t = x/V$, we obtain $\beta = c/V$. This is the limit value for passive causality violation; for $\beta > c/V$ we will have $t' < 0$.

If we write the signal velocity V as $V = c(1+\varepsilon)$, with $\varepsilon \ll 1$, ie V is only slightly superluminal, then the boost parameter β required to see the causality violation is $\beta \simeq 1-\varepsilon$. Qualitatively, if the signal velocity exceeds light velocity only by a small amount, then an observer must move almost at light velocity in order to be able to see a causality violation.

This elementary fact is already sufficient, in our opinion, to stimulate a reflection on the reality of causality violation in the case of signals which are only slightly superluminal. Consider, for instance, the superluminal neutrinos claimed by OPERA and then retracted, in the midst of a worldwide debate. The causality violation implied by their existence would only be observed in "laboratories" moving with a speed smaller than c by a few parts in one million. Is this a real logical evidence of causality violation?

One could correctly argue that physics has often made major progress by insisting on some fundamental symmetry or conservation law. Furthermore, the entire theory of Special Relativity is based on Gedanken-experiments which are difficult to implement in practice, and yet its logical conclusions are correct. "We do not need", one could say, "to really be able to move at almost-light velocity in order to conclude that a causality violation occurs". In other cases, however, a progress has been obtained in fundamental physics by distinguishing between quantities which are really observable, and quantities which are only an abstraction.

A. Emission of a bunch of tachyons from a source

As a further elementary exercise, let us consider the emission of several tachyons from a source at the same instant. (Hypothetical unstable particles travelling with superluminal velocity are usually called "tachyons"). To help our intuition, let us make a simple graphical and numerical example in units such that $c = 1$. The Lorentz transformations have the form $t' = \gamma(t - \beta x)$, $x' = \gamma(x - \beta t)$. All tachyons are emitted from $A(0, 0)$. Consider an event $D(3, 1)$. This represents the detection of a tachyon of velocity $V = +3c$ emitted from the origin at the instant $t = 0$ and propagating in the positive x direction. In order to observe a passive violation we should pass to a reference system in relative motion in the positive x direction with rapidity $\beta > c/V$, i.e. in this case $\beta > 1/3$. Let us choose for instance $\beta = 1/2$, whence $\gamma = (1 - \beta^2)^{-1/2} = 2/\sqrt{3}$. We obtain as new coordinates of D approximately

$$\begin{aligned} x'_D &= \tfrac{2}{\sqrt{3}}\left(3 - \tfrac{1}{2}\cdot 1\right) \approx 2.9 \\ t'_D &= \tfrac{2}{\sqrt{3}}\left(1 - \tfrac{1}{2}\cdot 3\right) \approx -0.6 \end{aligned} \tag{1}$$

There is indeed passive violation (because t'_D is negative). Clearly if the detector is not placed at $x = 3$ but closer to the origin or farther from it, say $x = 3k$, with $k > 0$, then the detection event D_1 has coordinates $D_1(3k, k)$ and the transformed coordinates D'_1 are multiplied by k, without affecting the sign (in the figure, $k = 2$).

Therefore all tachyons emitted in the positive x direction in the laboratory system will appear, in the moving system, as emitted in D', D'_1 ... and absorbed in $A = A'$. Those which travel for longer are emitted earlier, in proportion to the travelled distance. In the moving frame the tachyons appear to have negative velocity (although the reference system is slower and is not "overtaking" them). So if an "explosion" occurs at the origin, with the emission of several tachyons, the moving observer will see them starting from various detectors at different times and converging in the origin at the same instant. If the moving observer believes, as it is reasonable, that the explosion is an irreversible process, then he should conclude that either (1) Not all frames are equivalent, or (2) Tachyons, if they exist, violate the principle of entropy increase, or (3) Local realism is not applicable to tachyons and it is not consistent to distinguish between source and target in a process involving a tachyon exchange (compare Sect. V).



FIG. 2: Hypothetical explosive process in A with emission of several tachyons with the same velocity, which are then detected in D, D_1, ... In a moving system with passive causality violation, the tachyons appear to be emitted in D', D_1' ... and absorbed in $A = A'$. They appear to start from various "detectors" at different times and to converge in the origin at the same instant.

III. ACTIVE CAUSALITY VIOLATION

Let us take one more logical step and look at the velocity requirements for active causality violation, in the form of the so-called "tachyon anti-telephone" first discussed by Tolman ([6–8] and ref.s). The argument is written down in detail below, also in order to generalize it later to the case of re-emission with delay. We admit the existence of some device which emits particles with propagation velocity $V > c$ ("tachyons"), and also the existence of efficient detectors for such particles. We want to use these particles for communication in spacetime. Let us also suppose, for a start, that the emission and detection lags can be disregarded, so that the communication timing is determined only by the propagation intervals. We ask if it is possible not only to observe a causality violation in some reference system, but to actively "send secondary superluminal signals back in time" and switch off the source of the primary signals even before their emission (Fig. 3). This is an even more impressive logical evidence of causality violation. What are the velocity requirements?

Consider two reference systems: the first one is the "laboratory system", with coordinates (x, t), the second one the "moving system", with coordinates (x', t'), which is moving with respect to the laboratory system in the positive x direction, with velocity $v = \beta c$. Suppose that the origins of the two systems coincide at the initial time, that is, the origin $(0, 0)$ denotes the same event in the two reference systems. In the laboratory system there is a tachyon emitter at the origin $x = 0$; at the time $t = 0$ this emits a primary tachyon which travels in the positive x direction and is detected at time t_B by a receiver placed at $x = x_B$. Therefore the event $A(0, 0) = A'(0, 0)$ is the emission of



FIG. 3: Regge-Tolman paradox: in the laboratory system a primary tachyon is emitted in A and detected in B. In a system moving with rapidity $\bar{\beta}$ with respect to the laboratory it is possible to emit a secondary tachyon (event B_1) which reaches the origin of the laboratory system (event C) at the same time of the primary emission. We suppose initially that B and B_1 coincide (no delay between the reception of the primary tachyon in B and the emission of the secondary tachyon in B_1). The velocity requirement for $\bar{\beta}$ is found to be given by eq. (10), namely $\bar{\beta} = 2Vc/(c^2 + V^2)$.

the particle, and $B(x_B, t_B)$ its reception. We have $t_B = x_B/V$, with $V > c$. In the moving system the coordinates of the reception are $B'(x'_B, t'_B)$ and are related to the coordinates x_B, t_B by the Lorentz transformation

$$\begin{aligned} x'_B &= \gamma x_B - \beta\gamma c t_B \\ t'_B &= \gamma t_B - \beta\gamma \tfrac{x_B}{c} \end{aligned} \tag{2}$$

Now suppose that just when the laboratory detector is hit by the “primary” tachyon, a moving emitter (which is at rest in the moving system) is near B and sees the detection with negligible delay. The moving emitter then emits a “secondary” tachyon, in the negative x' direction. We call the secondary emission Event B_1; it coincides with B in the absence of any emission delay, and for now let us suppose that this is the case.

The event C is the transit of the secondary tachyon near the origin of the laboratory system, where the primary emitter is placed. This event has coordinates $(0, t_C)$ in the laboratory system and (x'_C, t'_C) in the moving system. Finally suppose that in the moving system several detectors are placed along the path of the secondary tachyon; if one of these detectors is near the primary emitter and receives the secondary tachyon, then it is programmed to disable the primary emitter.

Our task is to check under which conditions the time of the event C, in the laboratory system, is positive, zero or negative. If $t_C > 0$, then there is no causality violation, because the switching-off of the primary emitter, ultimately caused by the primary emitter itself, occurs after the primary emission. On the contrary, if $t_C \leq 0$, then we are confronted with active causality violation. It is straightforward to prove, using diagrams representing the spacetime trajectories of the primary and secondary tachyons, that t_C can become negative if the velocity of the moving system is sufficiently close to c. Here we want to compute the exact "critical" value $\bar{\beta}$ of the β parameter for which $t_C = 0$. For $\beta > \bar{\beta}$ active causality violation occurs.

We denote with τ the time elapsed, in the moving system, since the event B_1. At the time $(t'_B + \tau)$, in the moving system, the secondary tachyon emitted in B_1 is at the position

$$x' = x'_B - V\tau \tag{3}$$

In the laboratory system this position is transformed to

$$x = \gamma x' + \beta\gamma c(t'_B + \tau) \tag{4}$$

Setting $x = 0$ and solving simultaneously (3) and (4), we find the time τ_C that the secondary tachyon takes (for a fixed β), to reach the point of the primary emission. The result is

$$\tau_C = \frac{x'_B + \beta c t'_B}{V - \beta c} \tag{5}$$

and the instant when this happens is

$$t'_C = t'_B + \tau_C \tag{6}$$

Now, by imposing that $t'_C = 0$, we find the critical value of β such that the event C (secondary tachyon arrives at the emission location of the primary tachyon) occurs at the same time as the primary emission, i.e. $(x'_C, t'_C) = (0, 0)$. (Note that we have set $x'_C = 0$ already, after eq. (4)). From (5), (6) we obtain

$$\frac{x'_B + \bar{\beta} c t'_B}{V - \bar{\beta} c} + t'_B = 0 \tag{7}$$

and hence

$$x'_B + V t'_B = 0 \tag{8}$$

Transforming into the laboratory system:

$$\bar{\gamma} x_B - \bar{\beta}\bar{\gamma} c t_B + V\left(\bar{\gamma} t_B - \bar{\beta}\bar{\gamma}\frac{x_B}{c}\right) = 0 \tag{9}$$

and finally, recalling that $x_B = V t_B$, we obtain the desidered velocity requirement (graph in Fig. 4):

$$\bar{\beta} = \frac{2V}{c + \frac{V^2}{c}} \tag{10}$$



FIG. 4: Velocity requirement for active causality violation (eq. (10)). The range of interest (superluminal signals) is $V/c > 1$.

For slightly superluminal signals ($V = c(1+\varepsilon)$) one finds $\bar{\beta} \simeq 1 - \varepsilon^2/2$. For instance, for the X-shaped waves of Mugnai et al. [2], superluminality of 7 % leads to $\bar{\beta} \simeq 99.8$ %, while superluminality of 25 % leads to $\bar{\beta} \simeq 97$ %.

IV. MODIFICATION OF THE VELOCITY REQUIREMENTS IN THE PRESENCE OF A DEAD TIME

Let us now suppose that the emission of the second tachyon is delayed with respect to the detection of the primary tachyon. Between the events B and B_1 there will be a certain dead time Δt, and eq. (3) must be modified as follows: when $\tau < \Delta t$ the second tachyon is not present, while when $\tau \geq \Delta t$ the position of the second tachyon is

$$x' = x'_B - V\left(t - \Delta t\right) \tag{11}$$

In the laboratory system this transforms to

$$x = \gamma x' + \beta\gamma c\left(t'_B + \tau\right) \tag{12}$$

Setting as before $x = 0$ and solving the system between (11) and (12) one obtains the time τ_C when the second tachyon arrives in C:

$$\tau_C = \frac{x'_B + \beta c t'_B + V\Delta t}{V - \beta c} \tag{13}$$

(This obviously reduces to (5) when $\Delta t = 0$.) Now we set $t'_C = t'_B + \tau_C = 0$ and find an equation for the critical value $\bar{\beta}$ necessary for active causality violation with delay. After some algebra we find the irrational equation

$$2 - \bar{\beta}\left(\frac{c}{V} + \frac{V}{c}\right) + \frac{\Delta t}{t_B}\sqrt{1 - \bar{\beta}^2} = 0 \tag{14}$$

(Again, note that this reduces to (10) for $\Delta t = 0$.) Define the following parameters, supposed to be known:

$$\begin{aligned} r &= \tfrac{c}{V} + \tfrac{V}{c} \\ s &= \tfrac{\Delta t}{t_B} \end{aligned} \tag{15}$$

The parameter r depends on the propagation velocity, while s depends on the ratio between the delay in the secondary emission and the flight time of the primary tachyon. (Note that this flight time and the distance from the emitter have no influence on the requirement for causality violation, when there is no delay.) In a typical situation where $V = c(1+\varepsilon)$ and the delay is small, one has $r \simeq 2 + \varepsilon^2$ and $s \ll 1$; but we shall also consider extreme cases of large superluminality ($r \gg 1$) and large delay or very small flight time ($s \gg 1$).

Eq. (14) is rewritten as

$$2 - \bar{\beta} r + s\sqrt{1 - \bar{\beta}^2} = 0 \tag{16}$$

First we show in an elementary way through series expansions that with a signal which is only slightly superluminal ($V = c(1+\varepsilon)$) we can compensate for any delay and we always obtain an active causality violation for some $\bar{\beta}$. In fact, substituting into (16) the Ansatz

$$\bar{\beta} = 1 - \frac{1}{2}\eta\varepsilon^2, \quad \text{with} \;\; 0 < \eta < 1 \tag{17}$$

we obtain

$$s = \frac{\varepsilon^2 (1-\eta) + g\left(\varepsilon^2\right)}{\varepsilon\sqrt{\eta}\left(1 - \frac{1}{8}\eta\varepsilon^2 + h\left(\eta\varepsilon^2\right)\right)} \approx \frac{\varepsilon(1-\eta)}{\sqrt{\eta}} \quad \text{for} \;\; \eta \to 0 \;\; \text{and } \varepsilon \text{ small} \tag{18}$$

(The functions g and h are at least quadratic.)

We see that by choosing η close to zero, we can obtain a value of s as large as we want. This means that, in reverse, it is always possible to find a solution for $\bar{\beta}$, no matter how large the delay $s = \Delta t / t_B$.

In order to find the general solution of (16) and examine it in the strongly superluminal case $V \gg c$, $r \gg 2$, we transform (16) into a quadratic equation. By taking the square on both sides, we find that the solution of this equation is also solution of the following equation

$$\bar{\beta}^2\left(r^2 + s^2\right) - 4\bar{\beta} r + 4 - s^2 = 0 \tag{19}$$

(There is also one spurious solution, see below.)

The discriminant $\Delta = 4s^2\left(s^2 + r^2 - 4\right)$ is always positive. The solutions of (19) are

$$\bar{\beta}_{1/2} = \frac{2r \pm s\sqrt{s^2 + r^2 - 4}}{r^2 + s^2} \tag{20}$$



FIG. 5: Velocity requirement for active violation in the fairly superluminal case with large delay ($r > 2$, s up to 25).

but it is straightforward to check that only the solution with the plus sign is acceptable as a solution of (16). The check is most readily done in the limit $s \ll r$. In this same limit, we can also rewrite (20) as

$$\bar\beta \approx \frac{2}{r} + \frac{s}{r} \quad (\text{for } s \ll r) \tag{21}$$

and we see directly that $0 < \bar\beta < 1$, as it should be. From eq. (21) we also see that when r is large and s is approximately 1 (delay approximately equal to propagation time), one has $\bar\beta \simeq 3/r$.

The fairly superluminal case with large delay ($r > 2$, $s \gg 1$) or strongly superluminal with large delay ($r \gg 2$, $s \gg 1$) are more difficult to study in an analytical way. By plotting $\bar\beta$ as a function of r and s (Fig. 5) one can check that the solution of (20) with the plus sign always gives a value of $\bar\beta$ in the range $0 < \bar\beta < 1$, and that $\bar\beta \to 1^-$ when $s \to +\infty$. This means that it is always possible to obtain an active causality violation, compensating any delay in the secondary emission, provided the moving emitter is travelling fast enough. (Having already shown this in eq. (18) in the case of slightly superluminal velocity, we should not be surprised to find a general confirmation for strongly superluminal signals.)

The conclusion above is mathematically clear and is a consequence of the singularity of the Lorentz transformation for $\beta \to 1^-$. Physically, however, it is paradoxical and shakes our faith in the connection between superluminal signals and active causality violation. Take, for instance, the following extreme case: we have a primary superluminal signal that travels for a small fraction of a second and hits a secondary emitter; a secondary tachyon is emitted with delay of, say, *one year*, and travels back to the primary emitter. There exist moving reference systems where the



FIG. 6: Trying to distinguish source (S) from target (T) in a tachyon exchange leads to inconsistencies. In (a), S is seen to emit a tachyon, while T absorbs it. The tachyon carries a positive energy and a positive momentum from S to T (big red arrow). S recoils to the left, T recoils to the right (black small arrows). (b) The same process as observed in a moving frame with passive causality violation. The tachyon appears to be emitted from T' and absorbed by S' at a later time (big red arrow). According to the Lorentz transformation (27) the tachyon carries a negative energy, thus the total energetic balance of the process is the same as observed in the rest system. The recoil momenta have the same sign as before, but the transferred momentum is still positive. Therefore the "direction" of the momentum exchange appears to be incompatible with the recoil, in the sense that after T' has emitted some positive momentum, it should not recoil to the right, and similarly for S'.

secondary tachyon still can reach the primary emitter before the primary emission!

As an interesting limit of eq. (20), let us evaluate the requirement for active violation in the case of large superluminality $r \gg 2$, supposing that the delay in the secondary emission is due to the propagation of a luminal precursor before the superluminal signal. Denoting $\xi = V/c$, we have in this case $r \simeq \xi$ and $s \simeq \xi$. One then finds $\bar{\beta} \simeq \sqrt{2}/2 \simeq 71\%$, independently from ξ. Therefore the necessity of a luminal precursor does not eliminate in principle the active violation (which is possible for any delay, as seen above), but makes it very difficult to obtain with a realistic device.

V. CONCEPTUAL PROBLEMS IN INTERACTIONS VIA TACHYON EXCHANGE

In this section we consider a process in which two sources interact through the exchange of tachyons. As mentioned in the Introduction, tachyons are unlikely to exist as real particles, but they can formally represent virtual particles which transmit an interaction. Yet some conceptual problems are encountered when one tries to describe the exchange interaction as a sequence of tachyon emission-propagation-absorption. In order to show this, we only need a few basic relations concerning kinematics and Lorentz transformations.

Extensions of Special Relativity exist, compatible with the relativity principle, which provide a complete framework for the kinematics of superluminal particles ([7] and ref.s). The usual definitions of energy and momentum are extended by introducing an imaginary mass $m = iM$:

$$E = \frac{mc^2}{\sqrt{1 - V^2/c^2}} = \frac{Mc^2}{\sqrt{V^2/c^2 - 1}} \tag{22}$$

$$\mathbf{p} = \frac{m\mathbf{V}}{\sqrt{1 - V^2/c^2}} = \frac{M\mathbf{V}}{\sqrt{V^2/c^2 - 1}} \tag{23}$$

Combining eq.s (22) and (23) one obtains

$$E^2 - p^2c^2 = m^2c^4 < 0 \tag{24}$$

$$\frac{\mathbf{p}}{E} = \frac{\mathbf{V}}{c^2} \tag{25}$$

Note that if $E < 0$ (tachyon observed from a system with passive violation, see below), then $M < 0$, and $\mathbf{V}$ and $\mathbf{p}$ are opposite.

The Lorentz transformations of the energy and momentum of the exchanged tachyon for a boost in the x direction are

$$\begin{aligned} E' &= \gamma \left(E - \beta c p_x\right) \\ p'_x &= \gamma \left(p_x - \beta \tfrac{E}{c}\right) \end{aligned} \tag{26}$$

Suppose that the momentum is directed along x. Remembering that $p_x/E = V_x/c^2$ and denoting the positive value of V_x simply as V, one has

$$\begin{aligned} E' &= \gamma E \left(1 - \beta \tfrac{V}{c}\right) \\ p'_x &= \gamma p_x \left(1 - \beta \tfrac{c}{V}\right) \end{aligned} \tag{27}$$

Suppose to be in the case of passive violation, i.e. to observe the tachyon from a moving reference frame with rapidity $\beta > c/V$ (Fig. 6). We see from (27) that $E' < 0$. This is consistent with the inversion of time in the tachyon propagation, as described in Sect. II, in the following sense: in the lab frame, the tachyon is seen to transfer a positive energy from the source to the target; in the moving frame we see the tachyon leave the target and carry away a negative energy which is delivered to the source at a later time; thus the final energetic balance is the same, in the sense that some energy has passed from the source to the target. Unfortunately, however, a similar reasoning does not hold for the momentum transfer, because from eq. (27) we see that in the moving frame $p' > 0$, like in the laboratory! (See proof in Fig. 7.) This property, though elementary, is quite puzzling in our opinion, and implies that it is inconsistent to say that in any circumstance two bodies interact through the exchange of tachyons "emitted" from one of the two bodies and



FIG. 7: Proof that conditions for passive violation and momentum inversion are incompatible. On the horizontal axis, ξ represents the ratio between tachyon velocity and light velocity. The range of interest for ξ is $\xi > 1$. On the vertical axis, β is the parameter of a Lorentz boost ($0 < \beta < 1$). Passive causality violation of the tachyon is observed for $\beta > c/V$ (Sect. I), therefore for $\beta > 1/\xi$, which is the region of the graph betwee the lines $\beta = 1/\xi$ and $\beta = 1$. From eq. (27) we see that the momentum of the tachyon, as observed from the moving system, changes sign if $1 - \beta c/V < 0$, i.e. if $\beta > \xi$. The latter is the region between the lines $\beta = \xi$ and $\beta = 1$, which does not have any point in common with the region of passive causality violation.

"absorbed" by the other. Actually, if two bodies interact through the exchange of tachyons, it is impossible to say which body emits them and which one absorbs them, and to say that at some time the tachyon is carrying energy E and momentum p from one body to the other. We can only consider the exchange process as a whole. (This should not be surprising if one recalls the failure of the local realism principle in typical quantum processes like the EPR phenomenon.)

It follows that if the conservation law of energy and momentum dictates a certain energy-momentum ratio either in the "source" or in the "target" in any of the reference frames, this constraint will apply even if (apparently) backwards in time. For instance, if a "source" emits a tachyon which is absorbed by a non-relativistic "target" particle T with energy-momentum ratio

$$\frac{E_T}{p_T} = \frac{\frac{1}{2}mv_T^2}{m_T v_T} = \frac{1}{2}v_T \tag{28}$$

the E/p ratio of the tachyon must be equal to $\frac{1}{2}v_T$ and therefore "the target determines the propagation velocity of the tachyon and the recoil of the source". (Again, this looks reasonable if one regards the interaction process as a whole quantum process.)

According to Recami et al. [7, 8], the Tolman paradox and other causal paradoxes proposed for tachyons can be solved on the basis of the Feynman-Stuckelberg"re-interpretation principle", which requires to regard the secondary tachyon emitted in the moving system as an anti-tachyon emitted in fact from the laboratory. Our reasoning is independent from the reinterpretation principle.

Acknowledgments

I am grateful to R.A. Lewis for useful discussions and for pointing out to me the velocity requirement for active violation.

[1] Steinberg, A.M., Kwiat, P.G. and Chiao, R.Y.: Measurement of the single-photon tunneling time. Phys. Rev. Lett. 71, 708-711 (1993) Enders, A. and Nimtz, G.: On superluminal barrier traversal. J. Phys. I France 2, 1693-1698 (1992) Mugnai D., Ranfagni A. and Ronchi L.: The question of tunneling time duration: A new experimental test at microwave scale. Phys. Lett. A 247, 281-286 (1998) Balcou P. and Dutriaux L.: Dual Optical Tunneling Times in Frustrated Total Internal Reflection. Phys. Rev. Lett. 78, 851-854 (1997)

[2] Mugnai D., Ranfagni A., Ruggeri R.: Observation of Superluminal Behaviors in Wave Propagation. Phys. Rev. Lett. 84, 4830-4833 (2000)

[3] Saari P., Reivelt K.: Evidence of X-Shaped Propagation-Invariant Localized Light Waves. Phys. Rev. Lett. 79: 4135-4138 (1997). Durnin J., Miceli J.J., Eberly J.H.: Diffraction-free beams. Phys. Rev. Lett. 58: 1499-1501 (1987)

[4] Adam T. et al. (The OPERA Collaboration): Measurement of the neutrino velocity with the OPERA detector in the CNGS beam. Available via arXiv. http://arxiv.org/abs/1109.4897. Cited March 22, 2013

[5] Longo, M.J.: Tests of relativity from SN1987A. Phys. Rev. D 36, 32763277 (1987)

[6] Benford, G., Book, D. and Newcomb, W.: The Tachyonic Antitelephone. Phys. Review D 2, 263-265 (1970)

[7] Recami E., Fontana F., Garavaglia R.: Special relativity and superluminal motions: a discussion of some recent experiments. Int. J. Mod. Phys. A, 15, 2793-2812 (2000). Recami E., Zamboni-Rached M., Nobrega K.Z., Dartora C.A., Hernandez-Figueroa H.E.: On the localized superluminal solutions to the Maxwell equations. IEEE J. of Sel. Topics in Quantum Electronics 9, 59-73 (2003)

[8] Recami E.: Tachyon Kinematics and Causality: a Systematic Thorough Analysis of the Tachyon Causal Paradoxes. Found. Phys. 17, 239-296 (1987)

[9] Toller M.: Maximal acceleration, maximal angular velocity, and causal influence. Int. J. Theor. Phys. 29, 963-984 (1990)

[10] Nesterenko V.V., Feoli A., Lambiase G. and Scarpetta G.: Regularizing property of the maximal acceleration principle in quantum field theory. Phys. Rev. D 60, 065001 (1999)